

# The mitochondrial genome of *Megascolia azurea* (Christ) (Hymenoptera: Scoliidae) with its phylogenetic implications

Zhen LIU<sup>①</sup>, Chun CHEN, Shiqi CHEN

Zoology Key Laboratory of Hunan Higher Education, Hunan University of Arts and Science, Changde, Hunan 415000, China

**Abstract:** We sequenced and annotated the mitochondrial genome of *Megascolia azurea* (Christ, 1791), the only representative of the genus *Megascolia* Betrem, 1964 in the family Scoliidae in China. It is a larval parasitoid of the rhinoceros beetles *Xylotrupes gideon* (Linnaeus, 1767) and *Oryctes rhinoceros* (Linnaeus, 1758) (Coleoptera). The mitogenome is 16,335 bp in length and possesses 38 mitochondrial genes. However, unlike most animal mitochondrial genomes, it contains one extra trnM gene. Gene rearrangement shows an inversion event involving protein-coding and tRNA genes. The protein-coding genes in this sequence and another 8 species from the Aculeata were used for phylogenetic analysis by maximum likelihood (ML) analysis and Bayesian inference (BI) analyses using *Cotesia vestalis* from the family Braconidae as the outgroup. The topology of the resulted phylogenetic tree demonstrated that *M. azurea* is sister to *Megacampsomeris prismatic*, a species also belonging to family Scoliidae. Scoliidae was recovered as a sister to the clade of (Apidae + (Chrysididae + (Tiphidae + Mutillidae) + (Formicidae + Vespidae))), followed by Pompilidae which was located at a basal position.

**Key words:** mitochondrial genome; phylogeny; scoliid wasp

## 天蓝巨土蜂的线粒体基因组及其系统发育含义（膜翅目：土蜂科）

刘珍<sup>①</sup>, 陈淳, 陈诗琦

湖南文理学院, 湖南省高校动物学重点实验室, 湖南 常德 415000

**摘要:** 天蓝巨土蜂 *Megascolia azurea* (Christ, 1791) 是中国土蜂科巨土蜂属 *Megascolia* Betrem, 1964 的唯一代表且其为鞘翅目姬独角仙 *Xylotrupes gideon* (Linnaeus, 1767) 和椰蛀犀金龟 *Oryctes rhinoceros* (Linnaeus, 1758) 的幼虫寄生蜂。我们测通定了天蓝巨土蜂的线粒体基因组, 其长度为 16,335 bp 且拥有 38 个线粒体基因。与其他动物线粒体相比, 该线粒体基因包含一个额外的 trnM 基因。基因重排涉及蛋白编码基因和 tRNA 基因的倒置。我们通过最大似然法和贝叶斯干扰两种方法以菜蛾盘绒茧蜂为外群对天蓝巨土蜂和其他 8 种针尾部昆虫的线粒体蛋白编码基因进行了系统发育分析, 所获得的系统树拓扑结构表明, 土蜂科是 (蜜蜂科+ (青蜂科+ (沟土蜂科+ 蚁蜂科) + (蚁科+ 胡蜂科))) 的姐妹群, 而蛛蜂科与以上构成姐妹群并位于系统发育树的基部。

**关键词:** 线粒体基因组; 系统发育; 土蜂

Accepted 23 March 2023. Published online 16 May 2023. Published 25 June 2023.

① Corresponding author, E-mail: qingniao8.27@163.com

## Introduction

Scoliidae is a family distributed worldwide, with about 560 recognized species in 43 genera belonging to two subfamilies: Proscoliinae and Scoliinae (Osten 2005). Species of Scoliidae are larval parasitoids of scarabaeoid beetles and pollinators of various plants (Liu *et al.* 2021a). *Megascolia azurea* (Christ, 1791) is among the largest in body size and widely distributed in the Oriental Region and probably bivoltine in Hong Kong (Liu *et al.* 2021b; Christopher & Christopher 2021). *Xylotrupes gideon* (Linnaeus, 1767) and *Oryctes rhinoceros* (Linnaeus, 1758) (Coleoptera) were reported as hosts of this species (Betrem 1928).

Phylogenetic relationships among major lineages of the Aculeata have been studied over the years (Branstetter *et al.* 2017; Brothers 1975, 1993; Pilgrim *et al.* 2008; Wilson *et al.* 2013; Zheng *et al.* 2021). Mitochondrial genomes are one of the most highly studied genomic systems for molecular evolution and phylogenetic inference (Cameron 2014). Mitochondrial genomes have been extensively sequenced in the Hymenoptera and provide useful genetic markers for phylogenetic inferences (Dowton *et al.* 2009a, b; Tang *et al.* 2018; Zheng *et al.* 2018; Zheng *et al.* 2021). However, compared to other groups of the Hymenoptera, there is a lack of representative mitochondrial genomes for several major lineages of the Aculeata, such as the Scoliidae and Tiphidae (Tang *et al.* 2018). There is only one identified scoliid species, namely *Megacampsomeris prismatica* (*Campsomeris prismatica*)(unverified) that was found to be sequenced in NCBI (<https://www.ncbi.nlm.nih.gov/>) before this study. In this study, we determined the mitochondrial genome of *M. azurea* by next-generation sequencing. A thorough description of its genome features and analysis of phylogenetic relationships within the major Aculeata lineages are also provided to illuminate evolutionary patterns across the stinging wasps.

## Material and methods

### Collection, Identification and DNA extraction

Materials were all collected by hand netting in Taiyang Mountain (29.17N, 111.71E), Changde, China in August 2020. The collected specimens were stored in 100% ethanol and stored at  $-80^{\circ}\text{C}$  before DNA extraction.

Genomic DNA was extracted from the legs of single specimens using a Ezup Column Animal Genomic DNA Purification Kit (Sangon Biotech, China) following the manufacturer's protocols. Extracted genomic DNA were qualified by NanoPhotometer® (IMPLEN, CA, USA) and Qubit 3.0 (Invitrogen, Life Technologies, Carlsbad, CA, USA) and a Nanodrop 2000c Spectrophotometer (Thermo Scientific, Wilmington, DE, USA). Voucher specimens are deposited in the Laboratory of Hunan University of Arts and Science, Changde, China.

### Genome sequencing, assembly, annotation and analysis

The extracted genomic DNA of the specimen was sheared into fragments of approximately 350 bp in length using an Ultrasonic Processor Covaris S220 (Covaris, Inc. MS, USA). High-throughput sequencing libraries were constructed using the Illumina TruSeq

DNA PCR-Free HT Kit and sequenced using an Illumina Novaseq6000 with the strategy of producing 150 bp paired-ends by the Annoroad Gene Tech. (Beijing) Co., Ltd. Quality of raw sequencing reads was checked by FastQC version 0.11.3 (Andrews 2010), and low-quality reads and sites were filtered by Trimmomatic version 3.2.57 (Bolger *et al.* 2014).

The target mitochondrial reads were filtered out using BLAST (BLASTn with E value:  $1 \times 10^{-5}$ ) against a reference data set containing Braconidae mitochondrial genomes via the FastqExtract script (Crampton Platt *et al.* 2015). The mitochondrial genome of *M. azurea* was assembled by IDBA\_UD version 1.1.3 (Peng *et al.* 2012) and SPAdes version 3.15.2 (Bankevich *et al.* 2012) with default parameters.

Annotation of the assembled genome was performed by using MITOS Web Server (Bernt *et al.* 2013). Start and stop codons of protein-coding genes (PCGs) were manually adjusted in Geneious Prime v11 by referencing the published mitogenomes of Microgastrinae. The online tRNAscan-SE service (<http://lowelab.ucsc.edu/tRNAscan-SE/>) and XRNA tools were used to confirm locations of transfer RNA (tRNA) genes. Characteristics of mitochondrial genomes of sequenced species in Scoliidae were analyzed, including gene arrangements and base compositions and codon usages of PCGs according to methods mentioned in Yuan *et al.* (2022). Gene rearrangements were analyzed by comparison with the putative ancestral type of *Drosophila melanogaster*.

Mitochondrial genomes of 10 species from Hymenoptera were chosen to construct the phylogenetic relationships of Aculeata. *Cotesia vestalis* from Braconidae was chosen as the outgroup. The PCGs were aligned using G-INS-i algorithms implemented in MAFFT v7.407 (Katoh & Standley 2013). The best partition schemes of substitution models for the matrix were searched in PartitionFinder v2.1.12 with model selection = BIC and Branch lengths = unlinked between different subsets (Lanfear *et al.* 2012). MrBayes v3.2.7a (Ronquist & Huelsenbeck 2003) and IQtree v1.6.12 (Nguyen *et al.* 2015) were used to reconstruct the phylogenetic tree, respectively, based on nucleotide (NU) sequences of PCGs.

## Results

### Genome structure and organization

A total of 26 Gb filtered clean data were produced in this sequencing. The mitochondrial genome of *M. azurea* is 16,335 bp in length (GenBank accession OP661167), containing 13 protein-coding genes (PCGs), 23 transfer RNA genes (tRNAs), and two ribosomal RNA (rRNA) genes. However, unlike most animal mitochondrial genomes, it contains one extra trnM gene that can be detected using either the IDBA or SPAdes assemblers.

The overlapping nucleotides from seven adjacent genes in the mitochondrial genome of *M. azurea* were discovered to be up to 43 bp in total, as the maximum length of overlap was 25 bp in length, locating at one junction (rrnL-trnL1). The total intergenic nucleotides were 660 bp in length and dispersed between 27 adjacent genes, with the longest gene spacer, 127 bp in length dispersing between trnH and trnN. Neither overlapping nor space nucleotides were found between the other 6 adjacent genes. The majority strand (J-strand) encoded seven PCGs (atp6, atp8, cob, cox3, nad3, nad5, and nad6) and ten tRNAs (trnS2, trnT, trnH, trnN, trnR, trnA, trnG, trnD, trnC and trnY) while the remaining genes were encoded on the minority strand (N-strand).

**Table 1. Codon usage in the mitochondrial genome of *Megascolia azurea***

Codon	Number	RSCU	Amino acid	Codon	Number	RSCU
UUU	451	1.72	F	UUC	73	0.28
UUA	366	3.77	L	UUG	79	0.81
CUU	59	0.61		CUC	17	0.17
CUA	50	0.51		CUG	12	0.12
AUU	451	1.78	I	AUC	55	0.22
AUA	362	1.72		AUG	58	0.28
GUU	54	1.76	V	GUC	4	0.13
GUA	57	1.85		GUG	8	0.26
UCU	75	1.62	S	UCC	23	0.50
UCA	64	1.38		UCG	8	0.17
CCU	44	2.59	P	CCC	7	0.41
CCA	16	0.94		CCG	1	0.06
ACU	52	1.66	T	ACC	18	0.58
ACA	48	1.54		ACG	7	0.22
GCU	17	1.70	A	GCC	3	0.30
GCA	19	1.90		GCG	1	0.10
UAU	391	1.83	Y	UAC	37	0.17
UAA	443	1.74	*	UAG	65	0.26
CAU	54	1.89	H	CAC	3	0.11
CAA	63	1.64	Q	CAG	14	0.36
AAU	488	1.85	N	AAC	40	0.15
AAA	438	1.77	K	AAG	57	0.23
GAU	82	1.78	D	GAC	10	0.22
GAA	103	1.72	E	GAG	17	0.28
UGU	51	1.52	C	UGC	16	0.48
UGA	77	1.62	W	UGG	18	0.38
CGU	13	1.53	R	CGC	1	0.12
CGA	19	2.24		CGG	1	0.12
AGU	62	1.34	S	AGC	22	0.48
AGA	82	1.77		AGG	35	0.75
GGU	19	1.13	G	GGC	1	0.06
GGA	41	2.45		GGG	6	0.36

Abbreviation: RSCU, relative synonymous codon usage.

A relatively high A+T content in the mitochondrial genome is common in Hymenoptera (Oliveira *et al.* 2008). The mitochondrial genome of *M. azurea* contains a significant bias toward A/T in nucleotide composition by 85%, which ranges from 78.4% (cox1) to 91.9%

(nad2) with 42.7% of A, 8.1% of G, 42.3% of T, and 6.9% of C (Fig. 1). This high A+T content is consistent with the other scoliids (Zheng *et al.* 2021), showing Scoliidae is among the highest in A+T content in the Aculeata.




Figure 1. Base composition of PCGs and two rRNAs in the mitochondrial genome of *Megascolia azurea*.

The values of the GC skew in most genes were positive, except four genes (-0.14 in cox2, -0.08 in cox3, -0.26 in nad2 and -0.05 in nad5) and the value of the AT skew were also mostly positive, except for six genes (-0.15 in cox1 and nad1, -0.06 in cox2, and -0.11 in nad2, -0.12 in nad4 and -0.17 in nad4l) (Fig. 1).

The RSCUs in the mitochondrial genomes of *M. azurea* (Table 1) show an overwhelming preference toward the usage of A and T, especially at the third position, owing to genetic code degeneracies, which to some extent explains the high bias of A+T content. The total number of codons in the *M. azurea* mitochondrial genome is 5328. Among the available codons, there are five most widely used amino acids with their corresponding codons in the following order, Leu-UUA, Pro-CCU, Gly-GGA, Arg-CGA, and Ala-GCA, respectively.

#### Protein coding genes (PCGs)

The total length of 13 encoded PCGs is 11,011 bp in *M. azurea*, comprising 67% of the entire mitochondrial genome length. The size of the PCGs in the *M. azurea* mitochondrial genome is similar to the other corresponding ortholog (MH748671, 10,699 bp) in NCBI. The overall AT content of PCGs is 85.4%. The gene with the highest A + T content is nad2 with 91.9%, followed by atp8 with 91.5%.

All PCGs in the *M. azurea* mitochondrial genome initiated with the typical start codon ATN (seven with ATA, four with ATT, and two with ATG) as is the case of other Hymenoptera insects (Cameron *et al.* 2008; Castro & Dowton 2005; Cha *et al.* 2007; Crozier & Crozier 1993; Oliveira *et al.* 2008; Wei *et al.* 2009; Yuan *et al.* 2021). Approximately all

PCGs in the *M. azurea* mitochondrial genome terminate with a complete stop codon TAA or TAG. Only one PCG (*nad4*) used the incomplete stop codon T, which has been commonly reported in other invertebrates (Ojala *et al.* 1981).

### tRNA and rRNA genes




Figure 2. Predicted secondary structures for the 23 typical tRNA genes of *Megascolia azurea* mitochondrial genome. Base-pairing is indicated as follows: Watson–Crick pairs by lines, wobble GU pairs by dots and other non-canonical pairs by circles, tRNA, transfer RNA.

The entire 22 ancestral tRNAs commonly existing in the animal mitochondrial genome were successfully detected in the mitochondrial genome of *M. azurea*. But unlike other animal mitochondrial genomes, there was an extra trnM detected in the mitochondrial genome of *M. azurea*. These tRNA genes were scattered throughout the mitochondrial genome. Generally, the lengths of tRNAs range from 59 (trnS1) to 71 bp (trnM2). All tRNA genes in the mitochondrial genome of *M. azurea* have typical clover-leaf secondary structures except trnS1 is lacking the D loop (Fig. 2). The DHU-stem and TΨC-stem of the mitochondrial tRNA in *M. azurea* varied in length. The size of its anticodon loop of all tRNAs was 7 bp, except for that in trnS2 (9 bp). The size of the amino-acid acceptor stem of all tRNAs was 7 bp. Such variations are normal in other insect mitochondrial tRNAs (Sheffield *et al.* 2008; Wei *et al.* 2010; Peng *et al.* 2017). A total of 11 mismatched base pairs were detected with 9 G-U pairs and 2 U-U pairs.

### Gene rearrangements

Gene rearrangements were evident when compared with the ancestral type of *Drosophila melanogaster*. The rearrangements include protein-coding genes and tRNA genes. Inverse transposition occurred in gene clusters *trnI* to *trnK*, *trnH*, *trnS1* and *trnE*. Transposition occurred in *trnF* and local inversion occurred in *nad5* (Fig. 3). The rearrangements were similar to two unnamed mitogenomes of Scoliidae in Zheng *et al.* (2021) except an extra trnM gene and shows the rapid evolution of this group compared to other lineages of Hymenoptera.



Figure 3. Gene rearrangement in the mitochondrial genome of *Megascolia azurea*.

### Phylogenetic analyses




Figure 4. Phylogenetic tree of maximum likelihood (ML) methods and Bayesian inference (BI) using matrixes of 13 PCGs in mitochondrial genomes of 9 representative lineages in Aculeata and outgroup *Cotesia vestalis*. The support values of the corresponding nodes are shown above (left is bootstraps of ML, right is posterior probability of BI).

**Table 2. Optimal partitioning scheme for mitogenome dataset determined by PartitionFinder**

Subset	Subset Partitions	Sites	Best Model
1	c1p3, cbp3, c3p3, c2p3	1354	GTR+G
2	a6p3, n3p3, a8p3, n2p3, n6p3	958	GTR+G
3	n5p1, n4p1, n1p1, n4lp1, n3p1	1477	GTR+I+G
4	n6p1, n2p1, a8p1	622	GTR+I+G
5	n1p2, n5p2, n4p2	1273	GTR+I+G
6	cbp1, c2p1, c3p1, a6p1	1060	GTR+I+G
7	n4lp2, a8p2, n2p2, n6p2	704	GTR+I+G
8	n4lp3, n5p3, n4p3, n1p3	1355	GTR+G
9	n3p2, c2p2, cbp2, a6p2, c3p2	1182	GTR+I+G
10	c1p2	508	GTR+G
11	c1p1	508	GTR+I+G

We conducted phylogenetic analyses on the concatenated nucleotide sequences (PCG123) of *M. azurea* and other 8 species from the Aculeata. *Cotesia vestalis* from Braconidae was chosen as the outgroup to validate the phylogenetic position of *M. azurea* within Aculeata. Optimal partitioning strategy and models were selected using PartitionFinder v2.1.12 (Table 2). The resulted 10 data matrices were subjected to BI and ML analyses. The BI and ML analyses generated the same topology (Fig. 4). Generally, the support values of the corresponding nodes in the phylogenetic tree of BI were relatively higher than in MI in this study, which indicates MrBayes performs more reliably than ML methods in mitochondrial phylogenomic analyses of this group, and corresponds with other results of Hymenoptera (Dowton *et al.* 2009a; Wei *et al.* 2014). In our analyses, *M. azurea* is sister to *M. prismatic*, a species also belonging to the family Scoliidae, with a solid confidence value (bootstraps = 100, posterior probability = 1), which is consistent with the morphological characters. Pompilidae was always located at basal positions of the Aculeata in both analyses. The monophyly of the superfamily Pompiloidea were not recovered in both analyses, with *Mutilla europaea* (Mutillidae) clustering with Tiphidae in another clade. The sister relationship between the Formicoidea and Vespoidea was recovered in both analyses with *Vespula flaviceps* clustering with *Atta texana*, the same as with other results based on nuclear genomic data (Branstetter *et al.* 2017; Johnson *et al.* 2013; Peters *et al.* 2017), mitochondrial data (Zheng *et al.* 2021) and molecular combining morphological data (Pilgrim *et al.* 2008). Chrysididae was sister to the group (Tiphidae + Mutillidae) and (Formicoidea + Vespoidea), followed by Apidae and Scoliidae in our analyses. Mutillidae always had the longest branch length based on BI and ML analyses, which is congruent with a former study (Zheng *et al.* 2021). More mitogenomes sequences of more taxa will be needed to study the phylogenetic relationships within Scoliidae and other stinging wasps. Considerable uncertainty remains regarding higher-level relationships within Aculeata.



## Acknowledgements

This study was supported by the National Natural Science Foundation of China (32100351) and Hunan Provincial Natural Science Foundation of China (2023JJ30434). We thank Prof. Xue-xin Chen (Zhejiang University, Hangzhou) and the reviewers for their valuable comments on this manuscript..

## References

- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. Available from: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/> (accessed March 1, 2021)
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA & Pevzner PA. 2012. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology*, 19(5): 455–477.
- Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsche G, Pütz J, Middendorf M & Stadler PF. 2013. MITOS: Improved de novo metazoan mitochondrial genome annotation. *Molecular Phylogenetics and Evolution*, 69(2): 313–319.
- Betrem JG. 1928. Monographie der Indo-Australischen Scoliiden mit zoogeographischen Betrachtungen. *Treubia*, 9(supplement): 1–388.
- Bolger AM, Lohse M & Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, 30: 2114–2120.
- Branstetter MG, Danforth BN, Pitts JP, Faircloth BC, Ward PS, Buffington ML, Gates MW, Kula RR & Brady SG. 2017. Phylogenomic insights into the evolution of stinging wasps and the origins of ants and bees. *Current Biology*, 27: 1019–1025.
- Brothers DJ. 1975. Phylogeny and classification of the aculeate Hymenoptera, with special reference to Mutillidae. *The University of Kansas Science Bulletin*, 50: 483–648.
- Brothers DJ. 1993. Phylogeny of Aculeata: Chrysoidea and Vespoidea (Hymenoptera). *Journal of Hymenoptera Research*, 2: 227–304.
- Cameron SL. 2014. Insect mitochondrial genomics: Implications for evolution and phylogeny. *Annual Review of Entomology*, 59: 95–117.
- Cameron SL, Dowton M, Castro LR, Ruberu K, Whiting MF, Austin AD, Diement K & Stevens J. 2008. Mitochondrial genome organization and phylogeny of two vespid wasps. *Genome*, 51: 800–808.
- Castro LR & Dowton M. 2005. The position of the Hymenoptera within the Holometabola as inferred from the mitochondrial genome of *Perga condei* (Hymenoptera: Symphyta: Pergidae). *Molecular Phylogenetics and Evolution*, 34: 469–479.
- Cha SY, Yoon HJ, Lee EM, Yoon MH, Hwang JS, Jin BR, Han YS & Kim I. 2007. The complete nucleotide sequence and gene organization of the mitochondrial genome of the bumblebee, *Bombus ignitus* (Hymenoptera: Apidae). *Gene*, 392: 206–220.
- Christopher T & Christopher B. 2021. A review of the digger wasps (Insecta: Hymenoptera: Scoliidae) of Hong Kong, with description of one new species and a key to known species. *European Journal of Taxonomy*, 786(1): 1–92.
- Crampton-Platt A, Timmermans MJ, Gimmel ML, Kutty SN, Cockerill TD, Vun Khen C & Vogler AP. 2015. Soup to tree: the phylogeny of beetles inferred by mitochondrial metagenomics of a Bornean rainforest sample. *Molecular Biology and Evolution*, 32: 2302–2316.

- Crozier RH & Crozier YC. 1993. The mitochondrial genome of the honeybee *Apis mellifera*: Complete sequence and genome organization. *Genetics*, 133: 97–117.
- Dowton M, Cameron SL, Austin AD & Whiting MF. 2009a. Phylogenetic approaches for the analysis of mitochondrial genome sequence data in the Hymenoptera – A lineage with both rapidly and slowly evolving mitochondrial genomes. *Molecular Phylogenetics and Evolution*, 52: 512–519.
- Dowton M, Cameron SL, Dowavic JI, Austin AD & Whiting MF. 2009b. Characterization of 67 mitochondrial tRNA gene rearrangements in the Hymenoptera suggests that mitochondrial tRNA gene position is selectively neutral. *Molecular Biology and Evolution*, 26: 1607–1617.
- Katoh K & Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, 30: 772–780.
- Johnson BR, Borowiec ML, Chiu JC, Lee EK, Atallah J & Ward PS. 2013. Phylogenomics resolves evolutionary relationships among ants, bees, and wasps. *Current Biology*, 23: 2058–2062.
- Lanfear R, Calcott B, Ho SY & Guindon S. 2012. PartitionFinder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution*, 29(6): 1695–1701.
- Liu Z, van Achterberg C, He JH & Chen XX. 2021b. A checklist of Scoliidae (Insecta: Hymenoptera) from China. *Zootaxa*, 4966(2): 101–126.
- Liu Z, Yang SJ, Wang YY, Peng YQ, Chen HY & Luo SX. 2021a. Tackling the taxonomic challenges in the family Scoliidae (Insecta, Hymenoptera) using an integrative approach: A case study from southern China. *Insects*, 12(10): e892.
- Nguyen LT, Schmidt HA, Von Haeseler A & Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution*, 32(1): 268–274.
- Ojala D, Montoya J & Attardi G. 1981. tRNA punctuation model of RNA processing in human mitochondria. *Nature*, 290: 470–474.
- Oliveira D, Raychoudhury R, Lavrov DV & Werren JH. 2008. Rapidly evolving mitochondrial genome and directional selection in mitochondrial genes in the parasitic wasp *Nasonia* (Hymenoptera: Pteromalidae). *Molecular Biology and Evolution*, 25: 2167–2180.
- Osten T. 2005. Checkliste der Dolchwespen der Welt (Insecta: Hymenoptera, Scoliidae). *Naturforschende Gesellschaft*, 220: 1–62.
- Peng Y, Chen B & Li TJ. 2017. Sequencing and analysis of the complete mitochondrial genome of 2017 *Parapolybia crocea* (Hymenoptera: Vespidae). *Acta Entomologica Sinica*, 60(4): 464–474
- Peng Y, Leung HCM, Yiu SM & Chin FYL. 2012. IDBA-UD: A de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics*, 28(11): 1420–1428.
- Peters RS, Krogmann L, Mayer C, Donath A, Gunkel S, Meusemann K, Kozlov A, Podsiadlowski L, Petersen M, Lanfear R, Diez PA, Heraty J, Kjer KM, Klopstein S, Meier R, Polidori C, Schmitt T, Liu S, Zhou X, Wappler T, Rust J, Misof B & Niehuis O. 2017. Evolutionary history of the Hymenoptera. *Current Biology*, 27: 1013–1018.
- Pilgrim EM, von Dohlen CD & Pitts JP. 2008. Molecular phylogenetics of Vespoidea indicate paraphyly of the superfamily and novel relationships of its component families and subfamilies. *Zoologica Scripta*, 37: 539–560.
- Ronquist F & Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, 19(12): 1572–1574.
- Sheffield NC, Song H, Cameron SL & Whiting MF. 2008. A comparative analysis of mitochondrial genomes in Coleoptera (Arthropoda: Insecta) and genome descriptions of six new beetles. *Molecular Biology and Evolution*, 25(11): 2499–2509.

- Tang P, Zhu JC, Zheng BY, Wei SJ, Sharkey M, Chen XX & Vogler AP. 2018. Mitochondrial phylogenomics of the Hymenoptera. *Molecular Phylogenetics and Evolution*, 131: 8–18.
- Wei SJ, Li Q, van Achterberg K & Chen XX. 2014. Two mitochondrial genomes from the families Bethyridae and Mutillidae: independent rearrangement of protein coding genes and higher-level phylogeny of the Hymenoptera. *Molecular Phylogenetics and Evolution*, 77: 1–10.
- Wei SJ, Shi M, He JH, Sharkey MJ & Chen XX. 2009. The complete mitochondrial genome of *Diadegma semiclausum* (Hymenoptera: Ichneumonidae) indicates extensive independent evolutionary events. *Genome*, 52: 308–319
- Wei SJ, Tang E, Zheng LH, Shi M & Chen XX. 2010. The complete mitochondrial genome of *Evania appendigaster* (Hymenoptera: Evaniidae) has low A+T content and a long intergenic spacer between atp8 and atp6. *Molecular Biology Reports*, 37(4): 1931–1942.
- Wilson JS, von Dohlen CD, Forister ML & Pitts JP. 2013. Family-level divergences in the stinging wasps (Hymenoptera: Aculeata), with correlations to angiosperm diversification. *Evolutionary Biology*, 40: 101–107.
- Yuan RZ, Zhou JJ, Shu XH, Ye XQ, Tang P & Chen XX. 2022. The mitochondrial genome of *Chelonus formosanus* (Hymenoptera: Braconidae) with novel gene orders and phylogenetic implications. *Archives of Insect Biochemistry and Physiology*, 111(1): e21870.
- Zheng BY, Cao LJ, Tang P, van Achterberg K, Hoffmann AA, Chen HY, Chen XX & Wei SJ. 2018. Gene arrangement and sequence of mitochondrial genomes yield insights into the phylogeny and evolution of bees and sphecid wasps (Hymenoptera: Apoidea). *Molecular Phylogenetics and Evolution*, 124: 1–9.
- Zheng XY, Cao LJ, Chen PY, Chen XX, van Achterberg K, Hoffmann AA, Liu JX & Wei SJ. 2021. Comparative mitogenomics and phylogenetics of the stinging wasps (Hymenoptera: Aculeata), *Molecular Phylogenetics and Evolution*, 159: 107–119.